

Chapter 9.0 Literature Cited

Adams, S. M., A. M. Brown, and R. W. Goede. 1993. A quantitative health assessment index for rapid evaluation of fish condition in the field. *Transactions of the American Fisheries Society* 122:63-73.

Adams, S. M., and M. G. Ryon. 1994. A comparison of health assessment approaches for evaluating the effects of contaminant-related stress on fish populations. *Journal of Aquatic Ecosystem Health* 3:15-25.

Adams, S. M., K. D. Ham, M. S. Greeley, R. F. LeHew, D. E. Hinton, and C. F. Saylor. 1996. Downstream gradients in bioindicator responses: point source contaminant effects on fish health. *Canadian Journal of Fisheries and Aquatic Sciences* 53:2177-2187.

Allan, J. D. 1995. Stream ecology: structure and function of running waters. Chapman and Hall, London, UK.

Allan, J. D. 2004. Landscapes and riverscapes: the influence of land use on stream ecosystems *Annual Review of Ecology, Evolution, and Systematics* 35:257–84.

Anderson, J. B., and W. T. J. Mason. 1968. A comparison of benthic macroinvertebrates collected by dredge and basket samplers. *Journal of the Water Pollution Control Federation* 40:252-259.

Andrus, C. 2000. Experiences from the field: alcove sampling in the Willamette River, Oregon. In D. H. Landers, S. Allen-Gil, J. Reynolds, L. Carstensen, J. Sharber, P. Bayley, R. M. Hughes, and C. Andrus. *New Perspectives in Electrofishing*, EPA/600/R-99-108. US Environmental Protection Agency, Office of Research and Development, Washington, DC.

Angermeier, P. L., and J. R. Karr. 1984. Relationships between woody debris and fish habitat in a small warmwater stream. *Transactions of the American Fisheries Society* 113(3):716-726.

Angermeier P. L., and J. R. Karr. 1986. Applying an index of biotic integrity based on stream fish communities: considerations in sampling and interpretation. *North American Journal of Fisheries Management* 6:418-429.

Angermeier P. L., and R. A. Smogor. 1995. Estimating number of species and relative abundances in stream-fish communities: effects of sampling effort and discontinuous spatial distributions. *Canadian Journal of Fisheries and Aquatic Sciences* 52:936-949.

Angradi, T. R. (editor). 2006. Environmental Monitoring and Assessment Program: Great River Ecosystems, Field Operations Manual. EPA/620/R-06/002. US Environmental Protection Agency, Washington, DC. (available at: <http://www.epa.gov/emap/greatriver/fom.html>).

APHA (American Public Health Association). 1998. Standard methods for the examination of water and wastewater. 20th edition. American Public Health Association, Washington, DC.

Araujo, F. G., L. Fichberg, B. Carvalho, T. Pinto, and G. Peixoto. 2003. A preliminary index of biotic integrity for monitoring the condition of the Rio Paraiba do Sul, Southeast Brazil. *Environmental Management* 32(4):516-526.

Archibald, R. E. M. 1972. Diversity in some South African diatom associations and its relation to water quality. *Water Research* 6:1229-1238.

Bahls, L. L. 1992. Periphyton bioassessment methods for Montana streams. MT Department of Health and Environmental Services. Water Quality Bureau, Helena, Montana.

Bailey, R. C., M. G. Kennedy, M. Z. Dervish, and R. M. Taylor. 1998. Biological assessment of freshwater ecosystems using a reference condition approach: comparing predicted and actual benthic invertebrate communities in Yukon streams. *Freshwater Biology* 39:765-774.

Barbour, M. T., and J. Gerritsen. 1996. Subsampling of benthic samples: a defense of the fixed-count method. *Journal of the North American Benthological Society* 15:386-391.

Barbour, M. T., J. Gerritsen, G. E. Griffith, R. Frydenborg, E. McCarron, J. S. White, and M. L. Bastian. 1996. A framework for biological criteria for Florida streams using benthic macroinvertebrates. *Journal of the North American Benthological Society* 15:185-211.

Barbour, M. T., J. Gerritsen, B. D. Snyder, and J. B. Stribling. 1999. Rapid bioassessment protocols for use in streams and wadeable rivers: periphyton, benthic macroinvertebrates and fish. 2nd edition. EPA 841-B-99-002. US Environmental Protection Agency, Office of Water, Washington, DC.

Barbour, M. T., J. B. Stribling, and J. R. Karr. 1995. Multimetric approach for establishing biocriteria and measuring biological condition. Pages 63-77 in W. S. Davis and T. P. Simon (editors). *Biological assessment and criteria. Tools for water resource planning and decision making*. Lewis Publishers, Boca Raton, Florida.

Barbour, M. T., J. B. Stribling, and P. F. M. Verdonschot. 2006. (in press). The multihabitat approach of USEPA's rapid bioassessment protocols: Benthic macroinvertebrates. (*Journal of the Spanish Association of Limnology*) *Limnetica* 25(3-4).

Barbour, M. T., and C. O. Yoder. 2004. Technical guidelines: technical elements of a bioassessment program. Unpublished manuscript.

Bartsch L. A., W. B. Richardson, and T. J. Naimo. 1998. Sampling benthic macroinvertebrates in a large floodplain river: considerations of study design, sample size, and cost. *Environmental Monitoring and Assessment* 52:425-439.

Bayley, P. B. 1989. Aquatic environments in the Amazon basin, with an analysis of carbon sources, fish production, and yield. Pages 399-408 in D. P. Dodge (editor). Proceedings of the International Large River Symposium (LARS). Canadian Special Publication of Fisheries and Aquatic Sciences 106. Canadian Department of Fisheries and Oceans, Ottawa, Canada.

Bayley, P. B. 1995. Understanding large river -floodplain ecosystems. BioScience 45(3):153-158.

Beak, T. W., T. C. Griffing, and A. G. Appleby. 1973. Use of artificial substrate samplers to assess water pollution. In J. Cairns, and K. L. Dickson (editors). Biological methods for the assessment of water quality. American Society for Testing and Materials Spec. Technical Publication 528.

Beaumont, W. R. C., G. Peirson, and M. J. Lee. 2006. Factors affecting the characteristics and propagation of voltage gradient fields from electric fishing anodes. Fisheries Management & Ecology 13(1):47-52.

Benke, A. C. 2001. Importance of flood regime to invertebrate habitat in an unregulated river-floodplain ecosystem. Journal of the North American Benthological Society. 20:225-240.

Bell, D. 1973. The Coming of Post-Industrial Society. Basic Books, New York.

Benke, A. C., R. L. Henry III, D. M. Gillespie, and R. J. Hunter. 1985. Importance of snag habitat for animal production in southeastern streams. Fisheries 10(5):8-12.

Benke, A. C., and J. L. Meyer. 1988. Structure and function of a blackwater river in the southeastern USA. Internationale Vereinigung für Theoretische und Angewandte Limnologie, Verhandlungen 23:1209-1218.

Berger, W., H. McCarty, and R. K. Smith. 1996. Environmental laboratory data evaluation. Genium Publishing Corp, Amsterdam, New York.

Berkman, H. E., C. F. Rabeni, and T. P. Boyle. 1986. Biomonitor of stream quality in agricultural areas: fish vs invertebrates. Environmental Management 10:413-419.

Blocksom, K. A., and J. E. Flotemersch. 2005. Comparison of macroinvertebrate sampling methods for non-wadeable streams. Environmental Monitoring and Assessment 102:243-262.

Boivin T. G., D. Coates, D. Werle, and J. Baird. 2005. Remote sensing applications for information generation and monitoring for large river basin fisheries in tropical developing countries. Paper and poster prepared for the Second International Symposium on the Management of Large Rivers for Fisheries (February 2003), Phnom Penh, Cambodia.

Bryce, S. A., J. M. Omernik, and D. P. Larsen. 1999. Ecoregions: a geographic framework to guide characterization and ecosystem management. Environmental Practice 1:141-155.

Cao, Y., and C. P. Hawkins. 2005. Simulating biological impairment to evaluate the accuracy of ecological indicators. *Journal of Applied Ecology* 42:954-965.

Cao, Y., D. P. Larsen, and R. M. Hughes. 2001. Evaluating sampling sufficiency in fish assemblage surveys -- a similarity-based approach. *Canadian Journal of Fisheries and Aquatic Sciences* 58:1782-1793.

Cao, Y., D. P. Larsen, R. M. Hughes, P. L. Angermeier, and T. M. Patton. 2002. Sampling effort affects multivariate comparisons of stream assemblages. *Journal of the North American Benthological Society* 21(4):701-714.

Carboneau, P. E., S. N. Lane, and N. E. Bergeron. 2004. Catchment-scale mapping of surface grain size in gravel bed rivers using airborne digital imagery. *Water Resources Research* 40, W07202, doi:10.1029/2003WR002759.

Carpenter, S., N. F. Caraco, D. L. Correll, R. W. Howarth, A. N. Sharpley, and V. H. Smith. 1998. Nonpoint pollution of surface waters with phosphorous and nitrogen. *Issues in Ecology*, No. 3.

Carter, J. L., and V. H. Resh. 2001. After site selection and before data analysis: sampling, sorting and laboratory procedures used in stream benthic macroinvertebrate monitoring by USA state agencies. *Journal of the North American Benthological Society* 20(4):658-682.

Caton, L. R. 1991. Improved subsampling methods for the EPA rapid bioassessment benthic protocols. *Bulletin of the North American Benthological Society* 8:317-319.

Clark, M. J. R., and P. H. Whitfield. 1994. Conflicting perspectives about detection limits and about the censoring of environmental data. *Water Resources Bulletin* 30:1063-1075.

Clarke, R. 2000. Uncertainty in estimates of biological quality based on RIVPACS. Pages 39–54 in J. F. Wright, D. W. Sutcliffe, and M. T. Furse (editors). *Assessing the biological quality of fresh waters: RIVPACS and other techniques*. Freshwater Biological Association, Ambleside, UK.

Clements, W. H. 1991. Characterization of stream benthic communities using substrate-filled trays: colonization, variability, and sampling selectivity. *Journal of Freshwater Ecology* 6:209-221.

Cochran, W. G. 1977. *Sampling Techniques*. 3rd Ed. John Wiley & Sons. New York.

Cormier, S. M., S. B. Norton, G. W. Suter, II, D. Altfater, and B. Counts. 2002. Determining the causes of impairment in the Little Scioto River, Ohio: part II. Characterization of causes. *Environmental Toxicology and Chemistry* 21:1125-1137.

Courtemanch, D. L. 1996. Commentary on the subsampling procedures used for rapid bioassessments. *Journal of the North American Benthological Society* 15:381-385.

Craft, J. A., J. A. Stanford, and M. Pusch. 2002. Microbial respiration within a floodplain aquifer of a large gravel-bed river. *Freshwater Biology* 47(2):251-261.

Davies, P. E. 2000. Development of a national river bioassessment system (AUSRIVAS) in Australia. Pages 113-124 in J. F. Wright, D. W. Sutcliffe, and M. T. Furse (editors). *Assessing the biological quality of freshwaters: RIVPACS and other techniques*. Freshwater Biological Association, Ambleside, Cumbria, UK.

Davies, S. B., and S. K. Jackson. 2006. The Biological Condition Gradient: A conceptual model for interpreting detrimental change in aquatic ecosystems. *Ecological Applications* (in press).

Davis, W. S., and T. P. Simon (editors). 1995. *Biological assessment and criteria: tools for water resource planning and decision making*. Lewis Publishers, Boca Raton, Florida.

Davis, W. S., B. D. Snyder, J. B. Stribling, and C. Stoughton. 1996. Summary of state biological assessment programs for streams and wadeable rivers. EPA 230-R-96-007. Office of Policy, Planning, and Evaluation, US Environmental Protection Agency. Washington, DC.

Descy, J. P. 1979. A new approach to water quality estimation using diatoms. *Nova Hedwigia* 64: 305-323.

De Shon, J. 1995. Development and application of the invertebrate community index (ICI). Pages 217-244 in W. S. Davis, and T. P. Simon (editors). *Biological assessment and criteria*. Lewis Publishers, Boca Raton, Florida.

Dettmers, J. M., S. Gutreuter, D. H. Wahl, and D. A. Soluk. 2001. Patterns and abundance in main channels of the upper Mississippi River system. *Canadian Journal of Fisheries and Aquatic Sciences* 58:933-942.

Diamond, J. M., M. T. Barbour, and J. B. Stribling. 1996. Characterizing and comparing bioassessment methods and their results: A perspective. *Journal of the North American Benthological Society* 15:713-727.

Diamond, J. M., J. C. Hall, D. M. Pattie, and D. Gruber. 1994. Use of an integrated monitoring approach to determine site-specific effluent limits. *Water Environment Research* 66:733-743

Dolloff, C. A., and M. L. Warren. 2003. Fish relationships with wood in small streams. *Proceedings of the International Conference on Wood in World Rivers*. American Fisheries Society Special Symposium 37:179-193.

Dumont, S. C. and J. A. Dennis. 1997. Comparison of day and night electrofishing in Texas reservoirs. *North American Journal of Fisheries Management* 17:939-946.

Dunne, T., and L. B. Leopold. 1978. Water in environmental planning. W. H. Freeman and Co., New York.

Dynesius, M., and C. Nilsson. 1994. Fragmentation and flow regulation of river systems in the northern third of the world. *Science* 266:753-762.

Eberhardt, L. L. 1976. Quantitative ecology and impact assessment. *Journal of Environmental Management* 4:27-70.

Edsall, T. A., T. E. Behrendt, G. Cholwek, J. W. Frey, G. W. Kennedy, and S. B. Smith. 1997. Use of remote sensing techniques to survey the physical habitat of large rivers. U.S. Geological Survey Great Lakes Science Center. Contribution 983.

Ellis, J. L. and D. C. Schneider. 1997. Evaluation of a gradient sampling design for environmental impact assessment. *Environmental Monitoring and Assessment* 48:157-172.

Emery, E. B., T. P. Simon, F. H. McCormick, P. L. Angermeier, J. E. DeShon, C. O. Yoder, R. E. Sanders, W. D. Pearson, G. D. Hickman, R. J. Reash, and J. A. Thomas. 2003. Development of a multimetric index for assessing the biological condition of the Ohio River. *Transactions of the American Fisheries Society* 132:791-808.

Emery, E. B., J. A. Thomas, M. Bagley, and T. R. Angradi. 2006. Fish. In T. R. Angradi (editor). Environmental monitoring and assessment program: great river ecosystems field operations manual. EPA 620-R-06-002. US Environmental Protection Agency, Washington, DC.

Faith, D. P., C. L. Humphrey, and P. L. Dostine. 1991. Statistical power and BACI designs in biological monitoring: comparative evaluation of measures of community dissimilarity based on benthic macroinvertebrate communities in Rockdale Mine Creek, Northern Territories, Australia. *Australian Journal of Marine and Freshwater Research* 42:589-602.

Fausch, K. D., C. E. Torgersen, C. V. Baxter, and H. W. Li. 2002. Landscapes to riverscapes: bridging the gap between research and conservation of stream fishes. *BioScience* 52:483-498.

Faux, R. N., B. A. McIntosh, D. J. Norton, and J. R. Sedell. 1998. Thermal remote sensing of stream temperatures. Annual report, US Environmental Protection Agency, USDA Forest Service, and Bureau of Land Management.

Feminella, J. W. 2000. Correspondence between stream macroinvertebrate assemblages and 4 ecoregions of the southeastern USA. *Journal of the North American Benthological Society* 19:442-461.

Ferraro, S. P., F. A. Cole, W. A. DeBen, and R. C. Schwartz, 1989. Power-cost efficiency of eight macrobenthic sampling schemes in Puget Sound, Washington. *Canadian Journal of Fisheries and Aquatic Sciences* 46:2157–2165.

Fitzpatrick, F. A., I. R. Waite, P. J. D'Arconte, M. R. Meador, M. A. Maupin, and M. E. Gurtz. 1998. Revised methods for characterizing stream habitat in the National Water-Quality Assessment Program. US Geological Survey Water-Resources Investigations Report 98-4052.

Fjerdingstad, E. 1950. The microflora of the River Molleaa with special reference to the relation of benthic algae to pollution. *Folia Limnologica Scandanavica* 5:1-123.

Fjerdingstad, E. 1965. Taxonomy and saprobic valency of benthic phytomicroorganisms. *Internationale Revue der gesamten Hydrobiologie* 50(4):475-604.

Flannagan, J. F., and D. M. Rosenberg. 1982. Types of artificial substrates used for sampling freshwater benthic macroinvertebrates. In J. J. Cairns (editor). *Artificial substrates*. Ann Arbor Science, Ann Arbor, Michigan.

Flotemersch, J. E., Autrey, B. C., and Cormier, S. M. 2001. Comparisons of boating and wading methods used to assess the status of flowing waters. EPA 600-R-00-108. US Environmental Protection Agency, Cincinnati, Ohio.

Flotemersch, J. E., and K. A. Blocksom. 2005. Electrofishing in boatable rivers: Does sampling design affect bioassessment metrics?. *Environmental Monitoring and Assessment* 102:263-283.

Flotemersch, J. E., K. A. Blocksom, J. J. Hutchens, and B. C. Autrey. 2004. Association among invertebrates and habitat indicators for large rivers in the Midwest: how sampling distance, point-sampling of habitat, and subsample size effect measures of large river macroinvertebrate assemblages. EPA-600-R-04-177. US Environmental Protection Agency, Cincinnati, Ohio.

Flotemersch, J. E., K. A. Blocksom, J. J. Hutchens, and B. C. Autrey. 2006. Development of a standardized large river bioassessment protocol (LR-BP) for macroinvertebrate assemblages. *River Research and Applications* 22:775-790.

Flotemersch, J. E., and D. C. Jackson. 2003. Seasonal foraging by channel catfish on terrestrially burrowing crayfish in a floodplain-river ecosystem. *Ecohydrology and Hydrobiology* 3(1):61-70.

Flotemersch, J. E., and D. C. Jackson. 2005. Importance of crayfish to channel catfish in a floodplain-river ecosystem, in L. H. Fredrickson, S. L. King and R. M. Kaminski (editor). *Ecology and management of bottomland hardwood systems: The state of our understanding*. University of Missouri-Columbia. Gaylord Memorial Laboratory Special Publication No. 10. Puxico, Missouri.

Fore, L. S. 2004. Development and testing of biomonitoring tools for macroinvertebrates in Florida streams. Prepared for the Florida Department of Environmental Protection, Tallahassee, Florida.

Fore, L. S., J. R. Karr, and L. L. Conquest. 1994. Statistical properties of an index of biological integrity used to evaluate water resources. *Canadian Journal of Fisheries and Aquatic Sciences* 51:1077-1087.

Fore, L. S., J. R. Karr, and R. W. Wisseman. 1996. Assessing invertebrate responses to human activities: evaluating alternative approaches. *Journal of the North American Benthological Society* 15:212-231.

Frey, D.G. 1977. Biological integrity of water—a historical approach. Pages 127-140 in R. K. Ballantine and L. J. Guarria (editors). *The integrity of water*. US Environmental Protection Agency, Washington, DC.

Frissell, C. A., W. J. Liss, C. E. Warren, and M. D. Hurley. 1986. A hierarchical framework for stream habitat classification, viewing streams in a watershed context. *Environmental Management* 10:199-214.

Furse, M. T., D. Moss, J. F. Wright, P. D. Armitage, and R. J. M. Gunn. 1984. The influence of seasonal and taxonomic factors on the ordination and classification of running-water sites in Great Britain and on the prediction of their macroinvertebrate communities. *Freshwater Biology* 14:257-280.

Galat, D. L., and A. G. Frazier. 1996. Overview of river-floodplain ecology in the upper Mississippi River basin. In J.A. Kelmelis (editor). *Science for floodplain management into the 21st century*. Volume 3. Scientific Assessment and Strategy Team, US Government Printing Office, Washington, DC.

Galat, D. L., L. H. Fredrickson, D. D. Humburg, K. J. Bataille, J. R. Bodie, J. Dhorenwend, G. T. Gelwicks, J. E. Havel, D. L. Helmers, J. B. Hooker, J. R. M. Jones. 1998. Flooding to restore connectivity of regulated, large-river wetlands. *BioScience* 48:721-733.

Gale, W. F., and J. D. Thompson. 1974. Aids to benthic sampling by scuba divers in rivers. *Limnology and Oceanography* 19:1004-1007.

Gale, W. F., and J. D. Thompson. 1975. A suction sampler for quantitatively sampling benthos on rocky substrates in rivers. *Transactions of the American Fisheries Society* 104:398-405.

Gammon, J. R. 1973. The effect of thermal inputs on the populations of fish and macroinvertebrates in the Wabash River. Water Resources Research Center Technical Report 32, Purdue University, West Lafayette, Indiana.

Gammon, J. R. 1976. The fish populations of the middle 340 km of the Wabash River. Water Resources Research Center Technical Report 86, Purdue University, West Lafayette, Indiana.

Gammon, J. R., and T. P. Simon. 2000. Variation in a great river index of biotic integrity over a 20-year period. *Hydrobiologia* 422/423:291–304.

Ganasan, V., and R. M. Hughes. 1998. Application of an index of biological integrity (IBI) to fish assemblages of the rivers Khan and Kshipra (Madhya Pradesh), India. *Freshwater Biology* 40: 367–383.

Gauch, H. G. 1982. Multivariate analysis in community ecology. Cambridge University Press, Cambridge, UK.

- Gerritsen, J., M. T. Barbour, and K. King. 2000a. Apples, oranges, and ecoregions: on determining pattern in aquatic assemblages. *Journal of the North American Benthological Society* 19:487-496.
- Gerritsen, J., Burton, J., and M. T. Barbour. 2000b. A stream condition index for West Virginia wadeable streams. Prepared by Tetra Tech, Inc. for the US Environmental Protection Agency. Region 3 Environmental Services Division, and Office of Science and Technology, Office of Water, US Environmental Protection Agency, Washington, DC.
- Gerritsen, J., B. Jessup, and E. Leppo. 2002. Response of stream invertebrates to multiple anthropogenic stressors. *Water Environment Research Foundation Technical Report* 96-IRM-2.
- Gibson, G. R., M. T. Barbour, J. B. Stribling, J. Gerritsen, and J. R. Karr. 1996. Biological criteria: technical guidance for streams and small rivers (revised edition). EPA 822-B-96-001. US Environmental Protection Agency, Office of Water, Washington, DC.
- Goede, R. W., and B. A. Barton. 1990. Organismic indices and an autopsy-based assessment as indicators of health and condition of fish. *American Fisheries Society Symposium* 8:93-108.
- Gorman, O. T., and J. R. Karr. 1978. Habitat structure and stream fish communities. *Ecology* 59(3):507-515.
- Grafe, C. S. (editor). 2002. Idaho river ecological assessment framework: an integrated approach. Idaho Department of Environmental Quality. Boise, Idaho.
- Graf, W. L. 2001. Damage control: restoring the physical integrity of America's rivers. *Annals of the Association of American Geographers* 91:1-27.
- Graham, S. P. 1986. Comparison of day versus night electrofishing efficiency on largemouth bass at O'Shaughnessy Reservoir. Service Note 579, Division of Wildlife, Ohio Department of Natural Resources, Columbus, Ohio.
- Greenberg L, P. Svendsen, and A. Harby. 1996. Availability of microhabitats and their use by brown trout (*Salmo trutta*) and grayling (*Thymallus thymallus*) in the River Vojman, Sweden. *Regulated Rivers: Research & Management* 12:287–303.
- Griffith, M.B., P.R. Kaufmann, A.T. Herlihy, and B.H. Hill. 2001. Analysis of macroinvertebrate assemblages in relationship to environmental gradients in Rocky Mountain streams. *Ecological Applications* 11: 489-505.
- Griffith, M.B., B.H. Hill, A.T. Herlihy, and P.R. Kaufmann. 2002. Multivariate analysis of periphyton assemblages in relation to environmental gradients in Colorado Rocky Mountain streams. *Journal of Phycology* 38: 38-95.

Griffith, M.B., P. Husby, R.K. Hall, P.R. Kaufmann, and B.H. Hill. 2003. Analysis of macroinvertebrate assemblages in relation to environmental gradients among lotic habitats of California's Central Valley. *Environmental Monitoring and Assessment* 82(3): 281-309.

Grown, J., A. King, and F. Betts. 1999. The snag bug: a new method for sampling macroinvertebrate communities on large woody debris. *Hydrobiologia* 405:67-77.

Gutreuter, S., R. Burkhardt, and K. Lubinski. 1995. Long term resource monitoring program procedures: fish monitoring. LTRMP 95-P002-1. 42 pp. + Appendixes A-J. National Biological Service, Environmental Management Technical Center, Onalaska, Wisconsin, July 1995.

Hauer, F. R., and V. H. Resh. 1996. Benthic macroinvertebrates. In F. R. Hauer, and G. A. Lamberti (editors). *Methods in stream ecology*. Academic Press, New York.

Hawkins, C. P., and M. R. Vinson. 2000. Weak correspondence between landscape classifications and stream invertebrate assemblages: implications for bioassessment. *Journal of the North American Benthological Society* 19:501-517.

Hawkins, C. P., R. H. Norris, J. N. Hogue, and J. W. Feminella. 2000a. Development and evaluation of predictive models for measuring the biological integrity of streams. *Ecological Applications* 10:1456-1477.

Hawkins, C. P., R. H. Norris, J. Gerritsen, R. M. Hughes, S. K. Jackson, R. K. Johnson, and R. J. Stevenson. 2000b. Evaluation of the use of landscape classifications for the prediction of freshwater biota: synthesis and recommendations. *Journal of the North American Benthological Society* 19:541-556.

Hayes, D. B., C. P. Ferreri, and W. W. Taylor. 1996. Active fish capture methods. Pages 193-220 in, B. R. Murphy and D. W. Willis (editors). *Fisheries techniques*. American Fisheries Society, Bethesda, Maryland.

Herlihy, A. T., and C. W. Hendricks. 2000. Water chemistry and microbiology. Chapter 5 in J. M. Lazorchak, B. H. Hill, D. K. Averill, D. V. Peck, and D. J. Klemm (editors). *Environmental monitoring and assessment program - surface waters: field operations and methods for measuring the ecological condition of non-wadeable rivers and streams*. US Environmental Protection Agency, Cincinnati, Ohio.

Herzog, D. P., V. A. Barko, J. S. Scheibe, R. A. Hrabik, and D. E. Ostendorf. 2005. Efficacy of a benthic trawl for sampling small-bodied fishes in large river systems. *North American Journal of Fisheries Management* 25:594-603.

Hester, F. E., and J. S. Dendy. 1962. A multiple-plate sampler for aquatic macro-invertebrates. *Transactions of the American Fisheries Society* 91:420-421.

Hickman, G. D., and T. A. McDonough. 1996. Assessing the reservoir fish assemblage index: A potential measure of reservoir quality. Pages 85-97 in D. DeVries (editors). Reservoir Symposium – Multidimensional approaches to reservoir fisheries management. Reservoir Committee, Southern Division, American Fisheries Society, Bethesda, Maryland.

Hill, B. H., and A. T. Herlihy. 2000. ‘Periphyton’ in J. M. Lazorchak, B. H. Hill, D. K. Averill, D. V. Peck, and D. J. Klemm (editors). Environmental monitoring and assessment program – surface waters: field operations and methods for measuring the ecological condition of non-wadeable rivers and streams. USEPA 620-R-00-007. US Environmental Protection Agency, Cincinnati, Ohio.

Hill, B. H., A. T. Herlihy, P. R. Kaufmann, S. J. Decelles, and M. A. Vander Borgh. 2003. Assessment of streams of the eastern United States using a periphyton index of biotic Integrity. Ecological Indicators 2: 325-338.

Hill, B. H., A. T. Herlihy, P. R. Kaufmann, R. J. Stevenson, F. H. McCormick, and C. Burch Johnson. 2000. Use of periphyton assemblage data as an index of biotic integrity. Journal of the North American Benthological Society 19(1):50-67.

Hill, M. O. 1979. TWINSPAN - a FORTRAN program for arranging multivariate data in an ordered two way table by classification of the individuals and the attributes. Department of Ecology and Systematics, Cornell University, Ithaca, New York.

Hill, T. D., and D. W. Willis. 1994. Influence of water conductivity on pulsed AC and pulsed DC electrofishing catch rates for largemouth bass. North American Journal of Fisheries Management 14:202-207.

Hillebrand, H., C. D. Durselen, D. Kirschtel, U. Pollingher, and T. Zohary. 1999. Biovolume calculation for pelagic and benthic microalgae. Journal of Phycology 35:403-424.

Hilsenhoff, W. L. 1987. An improved biotic index of organic stream pollution. Great Lakes Entomologist 20:31-39.

Hocutt, C. H. 1981. Fish as indicators of biological integrity. Fisheries 6(6):28-30.

Hoffman, J. L. R. 2003. Susquehanna River pilot study: large river assessment project. Publication 228 Susquehanna River Basin Commission, Harrisburg, Pennsylvania.

Hubert, W. 1996. Passive capture techniques. Pages 157-192 in B. Murphy and D. Willis (editors). Fish Techniques. 2nd edition. American Fisheries Society, Bethesda, Maryland.

Huet, M. 1959. Profiles and biology of western European streams as related to fish management. Transactions of the American Fisheries Society 88:155-163.

Hughes, R. M., and J. R. Gammon. 1987. Longitudinal changes in fish assemblages and water quality in the Willamette River, Oregon. *Transactions of the American Fisheries Society* 116:196–209.

Hughes, R. M., and A. T. Herlihy. 2006 (accepted). Electrofishing effort needed to estimate consistent IBI scores in raftable Oregon Rivers. *Transactions of the American Fisheries Society*.

Hughes, R. M., P. R. Kaufmann, A. T. Herlihy, S. S. Intelmann, S.C. Corbett, M. C. Arbogast, and R. C. Hjort. 2002. Electrofishing distance needed to estimate fish species richness in raftable Oregon rivers. *North American Journal of Fisheries Management* 22:1229-1240.

Hughes, R. M., P. R. Kaufmann, A. T. Herlihy, T. M. Kincaid, L. Reynolds, and D. P. Larsen. 1998. A process for developing and evaluating indices of fish assemblage integrity. *Canadian Journal of Fisheries and Aquatic Sciences* 55: 1618-1631.

Hughes, R. M., D. P. Larsen, and J. M. Omernik. 1986. Regional reference sites: a method for assessing stream potentials. *Environmental Management* 10:629-635.

Hugueny, B., Camara, S., Samoura, B., Magassouba, M. 1996. Applying an index of biotic integrity based on fish communities in a west African river. *Hydrobiologia* 331:71-78.

Humphries, P., J. E. Growsn, L. G. Serafina, J. H. Hawking, A. J. Chick, and P. S. Lake. 1998. Macroinvertebrate sampling methods for lowland Australian Rivers. *Hydrobiologia* 364:209-218.

Hurlbert, S. J. 1984. Pseudo-replication and the design of ecological field experiments. *Ecological Monographs* 54:187-211.

ITFM (Intergovernmental Task Force on Monitoring Water Quality). 1995. The strategy for improving water quality monitoring in the United States: Final report of the Intergovernmental Task Force on monitoring water quality. US Geological Survey, Reston, Virginia.

Jennings, M. J., L. S. Fore, and J. R. Karr. 1995. Biological monitoring of fish assemblages in Tennessee Valley reservoirs. *Regulated Rivers* 11:263-274.

Jessup, B., and J. Gerritsen. 2000. Development of a multimetric index for biological assessment of Idaho streams using benthic macroinvertebrates. Prepared for the Idaho Department of Environmental Quality, Boise, Idaho.

Jessup, B. K., J. B. Stribling, and J. Gerritsen. 2002. Further evaluation of the Wyoming stream integrity index. Prepared by Tetra Tech, Inc. for the US Environmental Protection Agency, Region 8 Wyoming Department of Environmental Quality, and US Environmental Protection Agency.

Johnson, B. R., K. A. Blocksom, J. E. Flotemersch, and B. Autrey. 2004 (draft). Development of large river bioassessment protocols (LR-BP) for macroinvertebrate assemblages, quality assurance project plan. US Environmental Protection Agency, National Exposure Research Laboratory, Cincinnati, Ohio.

Johnson, B. L., W. B. Richardson, and T. J. Naimo. 1995. Past, present, and future concepts in large river ecology. *BioScience* 45(3):134-141.

Jones, K. K., C. A. Simenstad, D. L. Higley, and D. L. Bottom. 1990. Community structure, distribution, and standing stock of benthos, epibenthos, and plankton in the Columbia River estuary. *Progress in Oceanography* 25(1):211-241.

Junk, W. J., P. B. Bayley, and R. E. Sparks. 1989. The flood pulse concept in river-floodplain systems, in Dodge, D. P. (editor) *Proceedings of the International Large River Symposium*, Canadian Special Publication for Fisheries and Aquatic Sciences 106:110-127.

Karr, J. R. 1981. Assessment of biotic integrity using fish communities. *Fisheries* 6(6):21-27.

Karr, J. R. 1991. Biological integrity: a long-neglected aspect of water resource management. *Ecological Applications* 1:66-84.

Karr, J. R., and E. W. Chu. 1999. Restoring life in running waters: better biological monitoring. Island Press, Washington, DC.

Karr, J. R., and D. R. Dudley. 1981. Ecological perspective on water quality goals. *Environmental Management* 5:55-68.

Karr, J. R., K. D. Fausch, P. L. Angermeier, P. R. Yant, and I. J. Schlosser. 1986. Assessing biological integrity in running waters: a method and its rationale, special publication 5. Illinois Natural History Survey, Champaign, Illinois.

Karr, J. R., R. C. Heidinger, and E. H. Helmer. 1985. Effects of chlorine and ammonia from wastewater treatment facilities on biotic integrity. *Journal of the Water Pollution Control Federation* 57:912-915.

Kaufmann, P. R. (editor). 1993. Physical habitat. Pages 59-69 in R. M. Hughes (editor). Stream indicator and design workshop. EPA 600-R-93-138. US Environmental Protection Agency, Corvallis, Oregon.

Kaufmann, P. R. 2000. Physical habitat characterization - non-wadeable rivers. Chapter 6 in J. M. Lazorchak, B. H. Hill, D. K. Averill, D. V. Peck, and D. J. Klemm (editors). Environmental monitoring and assessment program - surface waters: field operations and methods for measuring the ecological condition of non-wadeable rivers and streams. US Environmental Protection Agency, Cincinnati, Ohio.

Kaufmann, P. R., P. Levine, E. G. Robison, C. Seeliger, and D. V. Peck. 1999. Quantifying Physical Habitat in Wadeable Streams. EPA 620/R-99/003. US Environmental Protection Agency, Washington, D.C.

Kaufmann, R. R., and E. G. Robison. 1997. Physical habitat assessment. In J. L. Lazorchak, D. J. Flemm, and D. V. Peck (editors). Environmental monitoring and assessment program – surface waters: field operations and methods for measuring the ecological condition of wadeable streams. EPA 620-R-94-004F. US Environmental Protection Agency, Washington, DC.

Kedwards, T. J., S. J. Maud, and P. F. Chapman. 1999. Community-level analysis of ecotoxicological field studies: II. Replicated design studies. *Environmental Toxicology and Chemistry* 18:158-166.

Keith L. H. (editor). 1988. Principles of environmental sampling. ISBN 0-8412-1173-6. ACS professional reference book. American Chemical Society, Columbus, Ohio.

Kelly, M. 1988. Mining and the freshwater environment. Elsevier Applied Science, London, UK.

Kelly, M. G. 1998. Use of the trophic diatom index to monitor eutrophication in rivers. *Water Research* 32:236-242.

Kentucky DOW. 2002. Methods for assessing biological integrity of surface waters in Kentucky. Division of Water, Water Quality Branch, Natural Resource and Environmental Protection Cabinet, Frankfort, Kentucky.

Kerans, B. L., and J. R. Karr. 1994. A benthic index of biotic integrity (B-IBI) for rivers of the Tennessee Valley. *Ecological Applications* 4:768-785.

King, R. S., M. E. Baker, D. F. Whigham, D. E. Weller, T. E. Jordan, P. F. Kazyak, and M. K. Hurd. 2005. Spatial considerations for linking watershed land cover to ecological indicators in streams. *Ecological Applications* 15:137-153.

Klemm, D. J., K. A. Blocksom, F. A. Fulk, A. T. Herlihy, R. M. Hughes, P. R. Kaufmann, D. V. Peck, J. L. Stoddard, W. T. Thoeny, M. B. Griffith, and W. S. Davis. 2003. Development and evaluation of a macroinvertebrate biotic integrity index (MBII) for regionally assessing Mid-Atlantic Highlands streams. *Environmental Management* 31(5): 656-669.

Klemm, D. J., P. A. Lewis, F. Fulk, and J. M. Lazorchak. 1990. Macroinvertebrate field and laboratory methods for evaluating the biological integrity of surface waters. EPA 600-4-90-030. US Environmental Protection Agency, Environmental Monitoring Systems Laboratory, Cincinnati, Ohio.

Kolkwitz, R., and K. Marsson. 1908. Ökologie der pflanzlichen Saproben. *Berichte der Deutschen botanischen Gesellschaft* 26A:505-519.

Knutson, M. G., and E. E. Klass. 1998. Floodplain forest loss and changes in forest community composition in the upper Mississippi River: a wildlife habitat at risk. *Natural Areas Journal* 18:138-150.

Krammer, K., and H. Lange-Bertalot. 1986. *Bacillariophyceae, Part 2, Volume 1: Naviculaceae*. In H. Ettl, J. Gerloff, H. Heynig, and D. Mollenhauer (editors). *Freshwater Flora of Middle Europe*. Gustav Fischer Publisher, New York.

Krammer, K., and H. Lange-Bertalot. 1988. *Bacillariophyceae, Part 2, Volume 2: Bacillariaceae, Epithemiaceae, Suriellaceae*. In H. Ettl, J. Gerloff, H. Heynig, and D. Mollenhauer (editors). *Freshwater Flora of Middle Europe*. Gustav Fischer Publisher, New York.

Krammer, K., and H. Lange-Bertalot. 1991a. *Bacillariophyceae, Part 2, Volume 3: Centrales, Fragilariaeae, Eunotiaceae*. In H. Ettl, J. Gerloff, H. Heynig, and D. Mollenhauer (editors). *Freshwater Flora of Middle Europe*. Gustav Fischer Publisher, Stuttgart, Germany.

Krammer, K., and H. Lange-Bertalot. 1991b. *Bacillariophyceae, Part 2, Volume 4: Achnanthaceae, Critical Supplement to Navicula (Lineolatae) and Gomphonema, Complete List of Literature for Volumes 1-4*. In H. Ettl, G. Gartner, J. Gerloff, H. Heynig, and D. Mollenhauer (editors). *Freshwater Flora of Middle Europe*. Gustav Fischer Publisher, Stuttgart, Germany.

Larsen, D. P. 1997. Sample survey design issues for bioassessment of inland aquatic ecosystems. *Human and Ecological Risk Assessment* 3(6):979-991.

Lange-Bertalot, Horst. 1979. Pollution tolerance of diatoms as a criterion for water quality estimation. *Nova Hedwigia* 64:285-304.

Lazorchak, J. M., B. H. Hill, D. K. Averill, D. V. Peck, and D. J. Klemm (editors). 2000. Environmental monitoring and assessment program – surface waters: field operations and methods for measuring the ecological condition of non-wadeable rivers and streams. US Environmental Protection Agency, Cincinnati, Ohio.

Lazorchak, J. M., Klemm, D. J., and D. V. Peck (editors). 1998. Environmental Monitoring and Assessment Program -Surface Waters: Field Operations and Methods for Measuring the Ecological Condition of Wadeable Streams. EPA/620/R-94/004F. US Environmental Protection Agency, Washington, D.C.

Legendre, P., and L. Legendre. 1998. Numerical ecology. 2nd English edition. Elsevier, New York.

Legleiter, C. J., D. A. Roberts, W. A. Marcus, and M. A. Fonstad. 2004. Passive optical remote sensing of river channel morphology and instream habitat: physical basis and feasibility. *Remote Sensing of Environment* 93:493– 510.

Lehtinen, R. M., N. D. Mundahl, and J. C. Madelczyk. 1997. Autumn use of woody snags by fishes in backwater and channel border habitats of a large river. *Environmental Biology of Fishes* 49:7-19.

Leland, H. V., and S. V. Fend. 1998. Benthic invertebrate distributions in the San Joaquin River, California, in relation to physical and chemical factors. *Canadian Journal of Fisheries and Aquatic Sciences* 55:1051-1067.

Leonard, P. M., and D. J. Orth. 1986. Application and testing of an index of biotic integrity in small, coolwater streams. *Transactions of the American Fisheries Society* 115:401-414.

Leopold, L. B., Wolman, M. G., and J. P. Mill. 1964. *Fluvial processes in geomorphology*: W.H. Freeman Press. San Francisco, California.

Li, J., A. Herlihy, W. Gerth, P. Kaufmann, S. Gregory, S. Urquhart, and D. Larsen. 2001. Variability in stream macroinvertebrates at multiple spatial scales. *Freshwater Biology* 46:87-97.

Liebmann, H. 1962. *Handbuch der frischwasser- und abwasserbiologie*. R. Oldenbourg, München, Germany.

Lisle, T. 1986. Effects of woody debris on anadromous salmonid habitat, Prince of Wales Island, southeast Alaska. *North American Journal of Fisheries Management* 6:538-550.

Lorenz, C. M., G. M. Van Dijk, A. G. M. Van Hattum, and W. P. Cofino. 1997. Concepts in river ecology: implications for indicator development. *Regulated Rivers Research and Management* 13:501-516.

Lowe, R. L. 1974. Environmental requirements and pollution tolerance of freshwater diatoms. EPA-670-4-74-005. US Environmental Protection Agency, Cincinnati, Ohio.

Lowe, R. L., and Y. Pan. 1996. Benthic algal communities as biological monitors. In R. J. Stevenson, M. L. Bothwell, and R. L. Lowe (editors). *Algal ecology. Freshwater benthic ecosystems*. Academic Press, San Diego, California.

Lowe-McConnell, R. H. 1975. *Fish communities in tropical freshwaters. Their distribution, ecology, and evolution*. Longman, London, UK.

Lubinski, K. S. 1993. A conceptual model of the upper Mississippi River system ecosystem. EMTC 93-T001. US Fish and Wildlife Service, Environmental Management Technical Center, Onalaska, Wisconsin.

Lubinski, K. S., and C. H. Theiling. 1999. Ecological status and trends of the upper Mississippi River system 1998: a report of the long term resource monitoring program (April 1999). LTRMP 99-T001. US Geological Survey, Upper Midwest Environmental Sciences Center, La Crosse, Wisconsin.

Lymburner, L., P. Hairsine, J. Walker, and A. Held. 2004. Riparian ecosystems and buffers: multi-scale structure, function, and management. AWRA Summer Specialty Conference Proceedings (June 28-30, 2004). Olympic Valley, California.

Lyons, J. 1992a. Using the index of biotic integrity (IBI) to measure environmental quality in warmwater streams of Wisconsin. General Technical Report, NC-149. Forest Service, US Department of Agriculture, St. Paul, Minnesota.

Lyons, J. 1992b. The length of stream to sample with a towed electrofishing unit when fish species richness is estimated. North American Journal of Fisheries Management 12:198–203.

Lyons, J., R. R. Piette, and K. W. Niermeyer. 2001. Development, validation, and application of a fish-based index of biotic integrity for Wisconsin's large warmwater rivers. Transactions of the American Fisheries Society 130:1077–1094.

Lyons, J., L. Wang, and T. D. Simonson. 1996. Development and validation of an index of biotic integrity for coldwater streams in Wisconsin. North American Journal of Fisheries Management 16:241-256.

Manly, B. J. F. 1994. Multivariate statistical methods: a primer. 2nd edition. Chapman and Hall, London, UK.

Manly, B. F. J. 2001. Statistics for environmental science and management. Chapman and Hall/CRC, Boca Raton, Florida.

Marchant, R., L. Barmuta, and B. C. Chessman. 1995. A preliminary study of the ordination and classification of macroinvertebrate communities from running waters in Victoria, Australia. Australian Journal of Marine Freshwater Research 45:945-962.

Maret, T. R., and D. S. Ott. 2004. Assessment of fish assemblages and minimum sampling effort required to determine biotic integrity of large rivers in Southern Idaho, 2002. US Geological Survey Water-Resources Investigations Report 03 –4274.

Maryland Department of Natural Resources. 1999. State of the streams: 1995-1997 Maryland biological stream survey results. CBWP-MANTA-EA-99-6. Maryland Department of Natural Resources, Annapolis, Maryland.

Mason, W. T., Jr. 1998. Macrofaunal monitoring in the Lower St. Johns River, Florida. Environmental Monitoring and Assessment 50:101-130.

Mason, W. T., C. I. Weber, P. A. Lewis, and E. C. Julian. 1973. Factors affecting the performance of basket and multiplate macroinvertebrate samplers. Freshwater Biology 3:409-436.

Matsushita, Y., and M. Shida. 2001. Comparison of catches between selective and conventional otter trawl for a coastal fishery. Reviews in Fisheries Science 9:33-42.

Matthews, R. A., P. F. Kondratieff, and A. L. Buikema, 1980. A field verification of the use of the autotrophic index in monitoring stress effects. *Bulletin of Environmental Contamination and Toxicology* 25:226-233.

McCormick, F. H., and R. M. Hughes. 2000. Aquatic vertebrates. Pages 10-1–10-14 in J. M. Lazorchak, B. H. Hill, D. K. Averill, D. V. Peck, and D. J. Klemm (editors). Environmental monitoring and assessment program – surface waters: field operations and methods for measuring the ecological condition of non-wadeable rivers and streams. EPA 620-R-00-007. US Environmental Protection Agency, Cincinnati, Ohio.

McCormick, F. H., D. V. Peck, and D. P. Larsen. 2000. Comparison of geographic classification schemes for Mid-Atlantic stream fish assemblages. *Journal of the North American Benthological Society* 19:385-404.

McCune, B., and J. B. Grace. 2002. Analysis of ecological communities. MjM Software Design, Gleneden Beach, Oregon.

McDonough, T. A., and G. D. Hickman. 1999. Reservoir fish assemblage index development: A tool for assessing ecological health in Tennessee Valley Authority impoundments. Pages 523-540 in T. P. Simon (editor). Assessing the sustainability and biological integrity of water resources using fish communities. CRC Press, Boca Raton, Florida.

Meador, M. R., T. F. Cuffney, and M. E. Gurtz. 1993. Methods for sampling fish communities as part of the National Water Quality Assessment Program. USGS/OFR 93-104. US Geological Survey, Washington, DC.

Meador, M. R., C. R. Hupp, T. F. Cuffney, and M. E. Gurtz. 1993. Methods for characterizing stream habitat as part of the National Water-Quality Assessment Program. U.S. Geological Survey, Open-File Report 93-408. Raleigh, North Carolina.

Mebane, C. A., T. R. Maret, and R. M. Hughes. 2003. An index of biological integrity (IBI) for Pacific Northwest rivers. *Transactions of the American Fisheries Society* 132:239-261.

Merritt, R. W, J. D. Allan, K. W. Cummins, K. J. Wessell, J. O. Wilhelm. 2005. Qualitative biological and habitat protocols for Michigan's non-wadeable rivers. Michigan Department of Environmental Quality, Lansing, Michigan.

Merritt, R., and K. W. Cummins. 2006. An Introduction to the Aquatic Insects of North America. 4th edition. Kendall/Hunt Publishing Company, Dubuque, Iowa.

Merritt, R. W., K. W. Cummins, and V. H. Resh. 1996. Design of aquatic insect studies: collecting, sampling, and rearing procedures. In R. W. Merritt, and K. W. Cummins (editors). An introduction to the aquatic insects of North America. 3rd edition. Kendall/Hunt Publishing, Dubuque, Iowa.

Mertes, L. A. K. 2002. Remote sensing of riverine landscapes. *Freshwater Biology* 47:799– 816.

Mertes, L. A. K., M. O. Smith, and J. B. Adams. 1993. Estimating suspended sediment concentrations in surface waters of the Amazon River wetlands from Landsat images. *Remote Sensing of Environment* 43:281–301.

Messer, J. J., R. A. Linthurst, and W. S. Overton. 1991. An EPA program for monitoring ecological status and trends. *Environmental Monitoring and Assessment* 17:67-78.

Meyer W. B., B. L. Turner, II (editors). 1994. Changes in land-use and land-cover: a global perspective. Cambridge University Press, Cambridge, UK.

Meyer, J. L. 1990. A blackwater perspective on riverine ecosystems. *BioScience* 40(9):643-651.

Minns, C. K., V .W. Cairns, R .G. Randall, and J. E. Moore. 1994. An index of biotic integrity (IBI) for fish assemblages in the littoral zone of Great Lakes areas of concern. *Canadian Journal of Fisheries and Aquatic Sciences* 51: 1804-1822.

Minshall, G. W., R. C. Peterson, K. W. Cummins, T. L. Bott, J. R. Sedell, C. E. Cushing, and R. L. Vannote. 1983. Interbiome comparison of stream ecosystem dynamics. *Ecological Monographs* 51:1-25.

Mississippi DEQ (Mississippi Department of Environmental Quality). 2006. Development and Application of the Mississippi Benthic Index of Stream Quality (M-BISQ). June 30, 2003. Prepared by Tetra Tech, Inc., Owings Mills, Maryland for Mississippi Department of Environmental Quality, Office of Pollution Control, Jackson, Mississippi.

Montgomery, D. R. and J. M. Buffington. 1998. Channel responses, classification, and response. Pages 13-42 in R. J. Naiman and R. E. Bilby (editors). *River Ecology and Management*. Springer-Verlag, New York.

Morales, E. A., and M. Potapova. 2000. Third NAWQA workshop on harmonization of algal taxonomy, (May 2000). Patrick Center for Environmental Research, the Academy of Natural Sciences, Philadelphia, Pennsylvania.

Moss, D., M. T. Furse, J. F. Wright, and P. D. Armitage. 1987. The prediction of the macro-invertebrate fauna of unpolluted running-water sites in Great Britain using environmental data. *Freshwater Biology* 17:41-52.

Moulton, S. R. II, J. G. Kennen, R. M. Goldstein, and J. A. Hambrook. 2002. Revised protocols for sampling algal, invertebrate, and fish communities as part of the national water-quality assessment program. USGS Open-File Report 02-150. US Geological Survey, Reston, Virginia.

Mouw, J. E. B., J. A. Stanford, and P. B. Alaback. 2003. Influences of fluvial processes and hyporheic exchange on floodplain plant diversity and productivity. *Oikos* (submitted).

Muli, J. R. , and K. M. Mavuti. 2001. The benthic macrofauna community of Kenyan waters of Lake Victoria. *Hydrobiologia* 458:83-90.

Naiman, R. J. 1983. The annual pattern and spatial distribution of aquatic oxygen metabolism in boreal forest watersheds. *Ecological Monograph* 53:73-94.

Naiman, R. J. 1998. Biotic stream classification. Pages 97-119 in R. J. Naiman and R. E. Bilby (editors). *River Ecology and Management*. Springer-Verlag, New York.

Naiman, R. J. and R.E. Bilby (editors). 1998. *River Ecology and Management: Lessons from the Pacific Coastal Ecoregion*. Springer-Verlag, New York.

National Research Council. 1992. *Restoration of aquatic ecosystems*. National Academy Press, Washington, DC.

Norton, D. J., M. A. Flood, B. A. McIntosh, [and 14 others]. 1996. Modeling, monitoring, and restoring water quality and habitat in Pacific northwestern watersheds. Final report. Environmental Program, Government Applications Task Force, Central Intelligence Agency, Washington, DC.

Norton, S. B., S. M. Cormier, M. Smith, and R. C. Jones. 2000. Can biological assessment discriminate among types of stress? A case study for the eastern cornbelt plains ecoregion. *Environmental Toxicology and Chemistry* 19(4):1113-1119.

Novak, M. A., and R. W. Bode. 1992. Percent model affinity, a new measure of macroinvertebrate community composition. *Journal of the North American Benthological Society* 11(1):80-85.

Novotny, D. W. 1990. Electric fishing apparatus and electric fields in I. G. Cowx, I. G. and P. Lamarque (editors). *Fishing with electricity, applications in freshwater fisheries management*: Oxford, England, Fishing News Books, Blackwell Scientific Publications, Ltd.

Novotny, D. W., and G. R. Priegel. 1974. Electrifishing boats, improved designs, and operational guidelines to increase the effectiveness of boom shocker. Wisconsin Department of Natural Resources Technical Bulletin No. 73, Madison, Wisconsin.

NWQMC (National Water Quality Monitoring Council). 2006. Data elements for reporting water quality monitoring results for chemical, biological, toxicological, and microbiological analytes. Prepared by methods and data comparability board for the advisory committee on water information. Technical Report: NWQMC 05-03. US Geological Survey, Reston, Virginia.

Oberdorff, T., and R. M. Hughes. 1992. Modification of an index of biotic integrity based to characterize rivers in the Seine-Normandie Basin, France. *Hydrobiologia* 228:117-330.

Ohio EPA (Ohio Environmental Protection Agency). 1987. Biological criteria for the protection of aquatic life: volumes I-III. Ohio Environmental Protection Agency, Columbus, Ohio.

Ohio EPA (Ohio Environmental Protection Agency). 1989. Biological criteria for the protection of aquatic life. Vol. III. Standardized field sampling and laboratory methods for assessing fish sampling and macroinvertebrate communities. Ohio EPA, Division of Water Quality Monitoring and Assessment. Columbus, Ohio.

Ohio EPA (Ohio Environmental Protection Agency). 1999. Association between nutrients, habitat, and the aquatic biota in Ohio rivers and streams. Ohio Environmental Protection Agency Technical Bulletin MAS/1999-1-1. Monitoring and Assessment Section, Division of Surface Water, Ohio Environmental Protection Agency, Columbus, Ohio.

Omernik, J. M. 1995. Ecoregions: a spatial framework for environmental management. Pages 49-62 in W. S. Davis, and T. P. Simon (editors). Biological assessment and criteria. Lewis Publishers, Boca Raton, Florida.

Osenberg, C. W., P. J. Schmitt, S. J. Holbrook, K. E. Abu-Saba, and R. Flegal. 1994. Detection of environmental impacts: natural variability, effect size, and power analysis. Ecological Applications 4(1):16-30.

Overton, W. S., D. White, and D. L. Stevens, 1991. Design report for EMAP, the environmental monitoring and assessment program. EPA 600-3-91-053. Office of Research and Development, US Environmental Protection Agency, Washington, DC.

Paller, M. H. 1995. Relationships among number of fish species sampled, reach length surveyed, and sampling effort in South Carolina coastal plain streams. North American Journal of Fisheries Management 15:110-120.

Pan, Y., R. J. Stevenson, B. H. Hill, and A. T. Herlihy. 2000. Ecoregions and benthic diatom assemblages in Mid-Atlantic highlands streams, USA. Journal of the North American Benthological Society 19:518-540.

Patrick, R. 1968. Effect of invasive rate, species pool and size area on the structure of diatom community. Proceedings of the National Academy of Sciences 58:1323-1335.

Patrick, R. 1977. Ecology of freshwater diatoms and diatom communities. In D. Werner (editor). The biology of diatoms. Botanical Monographs Volume 13. Blackwell Blackwell Scientific Publications, Ltd.

Patrick, R., N. J. Hohn, J. Wallace. 1954. A new method for determining the pattern of the diatom flora. Notulae Naturae (Philadelphia) No. 259.

Patrick, R., and C. W. Reimer. 1966. The diatoms of the United States, exclusive of Alaska and Hawaii. The Academy of Natural Sciences of Philadelphia, Philadelphia, Pennsylvania.

Patrick, R., and C. W. Reimer. 1967. The diatoms of the United States. Volume 1, Monograph 13. The Academy of Natural Sciences of Philadelphia, Philadelphia, Pennsylvania.

Patrick, R., and C. W. Reimer. 1975. The diatoms of the United States, exclusive of Alaska and Hawaii. The Academy of Natural Sciences of Philadelphia, Philadelphia, Pennsylvania.

Patton, T. M., W. A. Hubert, F. J. Rahel, and K. G. Gerow. 2000. Effort needed to estimate species richness in small streams on the Great Plains in Wyoming. *North American Journal of Fisheries Management* 20:394–398.

Paul, M. J., J. Gerritsen, and E. Leppo. 2002. Multivariate predictive invertebrate bioassessment models for Illinois streams. Draft Technical Report. Prepared by Tetra Tech, Inc. for Illinois Environmental Protection Agency, Springfield, Illinois.

Peck, D. V., D. K. Averill, A. T. Herlihy, R. M. Hughes, P. R. Kaugmann, D. J. Klemm, J. M. Lazorchak, F. H. McCormick, S. A. Peterson, M. R. Cappaert, T. Magee, and P. A. Monaco. 2006 (in press). Environmental Monitoring and Assessment Program – Surface Waters Western Pilot Study: Field Operations Manual for Non-Wadeable Rivers and Streams. EPA 620/R-06/xxx, U.S. Environmental Protection Agency, Washington, DC.

Penczak, T., and R. H. Mann. 1993. A preliminary evaluation of the effect of human activity on the fish populations of the Pilica River, central Poland. *Polish Archives of Hydrobiology* 40(1):101-115.

Petts, G.E. 1996. Water allocation to protect river ecosystems. *Regulated Rivers: Research and Management*. 12: 353-365.

Pitlo, J. 1992. An evaluation of the effect of a change in commercial harvest regulations on the channel catfish populations inhabiting the upper Mississippi River. Project F-109-R Des Moines. Federal Aid in Sport Fish Restoration, Iowa Department of Natural Resources, Des Moines, Iowa.

Plafkin, J. L., M. T. Barbour, K. D. Porter, S. K. Gross, and R. M. Hughes. 1989. Rapid bioassessment protocols for use in streams and rivers: benthic macroinvertebrates and fish. EPA 440/4/89/001. US Environmental Protection Agency, Office of Water Regulations and Standards, Washington, DC.

Poff, N. L., and J. D. Allan. 1995. Functional organization in stream fish assemblages in relation to hydrologic variability. *Ecology* 76:606-627.

Poff, N. L., J. D. Allan, M. B. Bain, J. R. Karr, K. L. Prestegaard, B. D. Richter, R. E. Sparks, and J. C. Stromberg. 1997. The natural flow regime: a paradigm for river conservation and restoration. *BioScience* 47:769-784.

Poff, N. L., and J. V. Ward. 1990. Physical habitat template of lotic systems: recovery in the context of historical pattern of spatiotemporal heterogeneity. *Environmental Management* 14(5): 629-645.

Poole, G. C., J. A. Stanford, C. A. Frissell, and S. W. Running. 2002. Three-dimensional mapping of geomorphic controls on flood-plain hydrology and connectivity from aerial photos. *Geomorphology* 48(4):329-347.

Porter, S. G., T. F. Cuffney, M. E. Gurtz, and M. R. Meador. 1993. Methods for collecting algal samples as part of the national water-quality assessment program. Open-File Report 93-409. US Geological Survey.

Poulton, B. C., M. L. Wildhaber, C. S. Charbonneau, J. F. Fairchild, B. G. Mueller, and C. J. Schmitt. 2003. A longitudinal assessment of the aquatic macroinvertebrate community in the channelized lower Missouri River. *Environmental Monitoring and Assessment* 85:23-53.

Prescott, G. W. 1962. Algae of the western Great Lakes area. Wm. Brown, Dubuque, Iowa.

Rabeni, C. F., and E. K. Gibbs. 1978. Comparison of two methods used by divers for sampling benthic invertebrates in deep rivers. *Journal of Fisheries Research Board of Canada* 35:332-336.

Rankin, E. T. 1989. The qualitative habitat evaluation index (QHEI): rationale, methods, and application. Ohio EPA, Division of Water Quality Planning and Assessment, Columbus, Ohio.

Rathbun, S. L. 1999. Sampling design issues for section 305(b) water quality monitoring. Appendix G in *Savannah River Basin REMAP: a demonstration of the usefulness of probability sampling for the purpose of estimating ecological condition in state monitoring programs*. EPA 904-R-99-002. US Environmental Protection Agency Region 4, SESD, Athens, Georgia.

Raven, P. J., N. T. H. Holmes, F. H. Dawson, and M. Everard. 1998. Quality assessment using river habitat survey data. *Aquatic Conservation: Marine and Freshwater Ecosystems* 8:477-499.

Raven, P. J., N. T. H. Holmes, F. H. Dawson, P. J. A. Fox, M. Everard, I. R. Fozzard, and K. J. Romen. 1998. River habitat quality: the physical character of rivers and streams in the UK and Isle of Man. ISBN 1-873760-429. Environment Agency, Bristol, England.

Reash, R. J. 1999. Considerations for characterizing Midwestern large river habitats. Pages 463-474 in T. P. Simon (editor). *Assessing the sustainability and biological integrity of water resources using fish communities*. CRC Press. Boca Raton, Florida.

Reckhow, K. H., and W. Warren-Hicks. 1997. Biological criteria: technical guidance for survey design and statistical evaluation of biosurvey data. EPA 822-B-97-021. US Environmental Protection Agency, Office of Water, Washington, DC.

Reeves, G. H., L. E. Benda, K. M. Burnett, P. A. Bisson, and J. R. Sedell. 1995. A disturbance-based ecosystem approach to maintaining and restoring freshwater habitats of evolutionarily significant units of anadromous salmonids in the Pacific Northwest. Pages 334-349 in J. L. Nielsen (editor). *Evolution and the aquatic ecosystem: defining unique units in population conservation*. American Fisheries Society Symposium 17, Bethesda, Maryland.

- Reimer, G. 1991. The ecological importance of floodplains for fish at the river March (Austria). *Archives of Hydrobiology* 121:355-363.
- Reisner, M. 1986. Cadillac desert: the American west and its disappearing water. Penguin Books, New York.
- Resh, V., and D. Rosenberg. 1984. The ecology of aquatic insects. Praeger, New York.
- Reynolds, J. B. 1996. Electrofishing. Pages 221-253 in B. R. Murphy and D. W. Willis (editors). *Fisheries techniques*, 2nd edition. American Fisheries Society, Bethesda, Maryland.
- Reynolds, L. A. T. Herlihy, P. R. Kaufmann, S. V. Gregory, and R. M. Hughes. 2003. Electrofishing effort requirements for assessing species richness and biotic integrity in Western Oregon streams. *North American Journal of Fisheries Management* 23:450-461.
- Reynoldson, T. B. and J. F. Wright. 2000. The reference condition: problems and solutions. Pages 293-303 in J. F. Wright, D. W. Sutcliffe, and M. T. Furse (editors). *Assessing the biological quality of fresh waters: RIVPACS and other techniques*. Freshwater Biological Association, Ambleside, Cumbria, UK.
- Richards, C., L. B. Johnson, and G. E. Host. 1996. Landscape-scale influences on stream habitats and biota. *Canadian Journal of Fisheries and Aquatic Science* 53(Supplement 1):295-311.
- Richter, B. D., D. P. Braun, M. A. Mendelson, and L. L. Master. 1997. Threats to imperiled freshwater fauna. *Conservation Biology* 11:1081-1093.
- Rogers, S. J., and T. W. Owens. 1995. Long Term Resource Monitoring Program Procedures: Vegetation monitoring. National Biological Service, Environmental Management Technical Center, Onalaska, Wisconsin, July 1995. LTRMP 95-P002-3. 10 pp. + Appendixes A-E.
- Rosenberg, D. M., and V. H. Resh. 1982. The use of artificial substrates in the study of freshwater benthic macroinvertebrates. In J. J. Cairns (editor). *Artificial substrates*. Ann Arbor Science, Ann Arbor, Michigan.
- Rosenberg, D. M., and V. H. Resh (editors). 1993. *Freshwater biomonitoring and benthic macroinvertebrates*. Chapman and Hall, New York.
- Rosenzweig, M. 1995. Species diversity in space and time. Cambridge University Press. London, UK.
- Roth, N. E., J. D. Allan, and D. L. Erickson. 1996. Landscape influences on stream biotic integrity assessed at multiple spatial scales. *Landscape Ecology* 11(3):141-156.
- Round, F. E., R. M. Crawford, and D. G. Mann. 1990. *The diatoms: biology and morphology of the genera*. Cambridge University Press, Cambridge, UK.

Sabater, S., F. Sabater, and J. Armengol. 1988. Relationships between diatom assemblages and physicochemical variables in the River Ter (NE Spain). *Internationale Revue der Gesamten Hydrobiologie* 73:171-179.

Salsburg, D. S. 1985. The religion of statistics as practiced in medical journals. *The American Statistician* 39:220-223.

Salsburg, D. S. 1986. Statistics for toxicologists. New York, Marcel Dekker, Inc.

Sanders, R. E. 1991. Day versus night electrofishing catches from near-shore waters of the Ohio and Muskingum Rivers. *Ohio Journal of Science* 192:1-59.

Sanders, R. E., R. J. Miltner, C. O. Yoder, and E. T. Rankin. 1999. The use of external deformities, erosion, lesion, and tumors (DELT anomalies) in fish assemblages for characterizing aquatic resources: a case study of seven Ohio streams. Pages 225-248 *in* T. P. Simon (editor). Assessing the sustainability and biological integrity of water resources using fish communities. CRC Press, Boca Raton, Florida.

Schmitt, C. J., V. S. Blazer, G. M. Dethloff, D. E. Tillitt, T. S. Gross, W. L. Bryant, L. R. DeWeese, S. B. Smith, R. W. Goede, T. M. Bartish, and T. J. Kubiak. 1999. Biomonitoring of environmental status and trends (BEST) program: field procedures for assessing the exposure of fish to environmental contaminants. *Information and Technology Report 1999-0007*. Biological Resources Division, US Geological Survey.

Scott, M. T., and L. A. Nielsen. 1989. Young fish distribution in backwaters and main-channel borders of the Kanawha River, West Virginia. *Journal of Fish Biology* 35:21-27.

Sedell, J. R., J. E. Richey, and F. J. Swanson. 1989. The river continuum concept: a basis for the expected ecosystem behavior of very large rivers? *Canadian Special Publication of Fisheries and Aquatic Sciences* 106:110-127.

Sheaffer, W. A., and J. G. Nickum. 1986. Backwater areas as nursery habitats for fishes in Pool 13 of the upper Mississippi River. *Hydrobiologia* 136:131-140.

Sheehan, R. J., and J. L. Rasmussen. 1999. Large rivers. Pages 529–559 *in* C. Kohler, W. A. Hubert (editors.). Inland fisheries management in North America. 2nd edition American Fisheries Society. Bethesda, Maryland.

Shelton, L. R. 1994. Field guide for collecting and processing stream-water samples for the national water-quality assessment program. Open-File Report 94-455. US Geological Survey, Sacramento, California.

Shepard, J. 2001. The state of the world's rivers. Center for Global Environmental Education, St. Paul, Minnesota.

Simon, T. P. 1991. Development of index of biotic integrity expectations for the ecoregions of Indiana. I. Central Corn Belt Plain. EPA-905-9-91-025. US Environmental Protection Agency, Region V, Chicago, Illinois.

Simon, T. P. (editor). 1999. Assessing the sustainability and biological integrity of water resources using fish communities. CRC Press, Boca Raton, Florida.

Simon, T. P., and J. Lyons. 1995. Application of the index of biotic integrity to evaluate water resources integrity in freshwater ecosystems. Pages 245–262 in W. S. Davis and T. P. Simon (editors). Biological assessment and criteria: tools for water resource planning and decision making. Lewis Publishers, Boca Raton, Florida.

Simon, T. P., and R. E. Sanders. 1999. Applying an index of biotic integrity based on great river fish communities: considerations in sampling and interpretation. Pages 475–505 in T. P. Simon (editor). Assessing the sustainability and biological integrity of water resources using fish communities. CRC Press, Boca Raton, Florida.

Simonson, T. D., J. Lyons, and P. D. Kanehl. 1994. Quantifying fish habitat in streams-transect spacing, sample size, and a proposed framework. North American Journal of Fisheries Management 14:607-615.

Simpson, J. C., and R. H. Norris. 2000. Biological assessment of river quality: development of AUSRIVAS models and outputs. Pages 125-142 in J. F. Wright, D. W. Sutcliffe, and M. T. Furse (editors). Assessing the biological quality of freshwaters: RIVPACS and other techniques. Freshwater Biological Association, Ambleside, Cumbria, UK.

Slack, K. V., R. F. Ferreira, and R. C. Averett. 1986. Comparison of four artificial substrates and the ponar grab for benthic invertebrate collection. Water Resources Bulletin 22:237-248.

Sladecek, V. 1973. System of water quality from the biological point of view. Archiv für Hydrobiologie – Advances in Limnology 7:1-218.

Smith, E. P. 2002. BACI design. Pages 141-148 in A. H. El-Shaarawi and W. W. Piegorsch, (editors). Encyclopedia of Environmetrics. John Wiley & Sons, Ltd, Chichester, UK.

Smith, E. P., D. R. Orvos, and J. J. Cairns. 1992. Impact assessment using the before-after-control-impact (BACI) model: concerns and comments. Canadian Journal of Fisheries and Aquatic Sciences 50:627-637.

Smith, F., S. Kulkarni, L. E. Myers, and M. J. Messner. 1988. Evaluating and presenting quality assurance sampling data. Chapter 10, pages 157-168 in L. H. Keith (editor). Principles of environmental sampling. ISBN 0-8412-1173-6. ACS professional reference book. American Chemical Society, Columbus, Ohio.

Smith, R. A., G. E. Schwartz, and R. B. Alexander. 1997. Regional interpretation of water quality monitoring data. Water Resources Research 33:2781-2798.

- Smith, S. B., A. P. Donahue, R. J. Lipkin, V. S. Blazer, C. J. Schmitt, and R. W. Goede. 2002. Illustrated field guide for assessing external and internal anomalies in fish. Information and Technology Report, 2002-0007. US Geological Survey (available at: http://www.cerc.usgs.gov/pubs/center/pdfDocs/ITR_2002_0007.pdf).
- Snyder, D. E. 2003. Electrofishing and its harmful effects on fish. Information and Technology Report. USGS/BRD/ITR—2003-0002. US Government Printing Office, Denver, Colorado.
- Sokal, R. R., and F. J. Rohlf. 1995. Biometry. 3rd edition. W. H. Freeman and Co., New York.
- Sparks, R. E., P. B. Bayley, S. L. Kohler, and L. L. Osborne. 1990. Disturbance and recovery of large floodplain rivers. Environmental Management 14:699-709.
- Stalnaker, C. B., R. T. Milhous, and K. D. Bovee. 1989. Hydrology and hydraulics applied to fishery management in large rivers. Page 13 in D. P. Dodge (editor). Proceedings of the International Eastern Rivers and Mountains Network – Phase I: Large River Symposium. Canadian Special Publication of Fisheries and Aquatic Sciences 106.
- StatSoft. 1994. STATISTICA Mac version 4.1. StatSoft Inc., Tulsa, Oklahoma.
- Steedman, R. J. 1988. Modification and assessment of an index of biotic integrity to quantify stream quality in southern Ontario. Canadian Journal of Fisheries and Aquatic Sciences 45:492-501.
- Stevens, D. L. 1997. Variable density grid-based sampling designs for continuous spatial populations. Environmetrics 8:167-95.
- Stevens, D. L., and A. R. Olsen. 1991. Statistical issues in environmental monitoring and assessment. Draft document – prepared for publication in Proceedings of the Section on Statistics and the Environment, American Statistical Association, Alexandria, Virginia.
- Stevens, D. L., and A. R. Olsen. 1999. Spatially restricted surveys over time for aquatic resources. Journal of Agricultural, Biological, and Environmental Statistics 4:415-28.
- Stevenson, R. J. 1996. An introduction to algal ecology in freshwater benthic habitats. Pages 3-30 in R. J. Stevenson, M. L. Bothwell, R. L. Lowe (editors). Algal ecology. Academic Press, San Diego, California.
- Stevenson, R. J., and L. Bahls. 1999. Periphyton protocols. In M. T. Barbour, J. Gerritsen, B. D. Snyder, J. B. Stribling (editors). Rapid bioassessment protocols for use in streams and wadeable rivers: periphyton, benthic macroinvertebrates, and fish. 2nd edition. EPA 841-B-99-002. Office of Water, US Environmental Protection Agency, Washington, DC.
- Stevenson, R. J., and J. P. Smol. 2002. Use of algae in environmental assessments. In J. D. Wehr, and R. G. Sheath, eds. Freshwater Algae in North America: Classification and Ecology. Academic Press, San Diego, California.

Stewart, J. G., and V. A. Barko. 2005. New records of the crystal darter (*Crystallaria asprella*) in the middle Mississippi River. *American Midland Naturalist* 154:471-473.

Stewart-Oaten, A., and J. R. Bence. 2001. Temporal and spatial variation in environmental assessment. *Ecological Monographs* 71:305-339.

Stewart-Oaten, A., W. W. Murdoch, and K. R. Parker. 1986. Environmental impact assessment: pseudoreplication in time? *Ecology* 67:929-940.

Stoddard, J. L., D. V. Peck, S. G. Paulsen, J. Van Sickle, C. P. Hawkins, A. T. Herlihy, R. M. Hughes, P. R. Kaufmann, D. P. Larsen, G. Lomnický, A. R. Olsen, S. A. Peterson, P. L. Ringold, and T. R. Whittier. 2005. An Ecological Assessment of Western Streams and Rivers. EPA 620/R-05/005, US Environmental Protection Agency, Washington, DC.

Stoddard, J. L., D. P. Larsen, C. P. Hawkins, R. K. Johnson, and R. H. Norris. 2006. Setting expectations for the ecological condition of streams: The concept of reference condition. *Ecological Applications* 16(4):1267-1276.

Stokesbury, K. D. E., J. B. Bighy, and S. W. Ross. Selectivity and efficiency of two otter trawls used to assess estuarine fish and macroinvertebrate populations in North Carolina. *Estuaries* 22(4):882-888.

Strahler, A. N. 1957. Quantitative analysis of watershed geomorphology. *Transactions of the American Geophysical Union* 38:913-920.

Stribling, J. B., and S. R. Davie. 2005. Design of a long-term environmental monitoring program for the Lake Allatoona/Upper Etowah River watershed, Georgia. Prepared for the US Army Corps of Engineers (Mobile District) and the Lake Allatoona/Upper Etowah River watershed partnership. Mobile, Alabama. (Available from: <http://allatoona.sam.usace.army.mil/Documents.htm>)

Stribling, J. B., Jessup, B. K., White, J. S., Boward, D., and Hurd, M. 1998. Development of a Benthic Index of Biotic Integrity for Maryland Streams. CBWP-MANTA-EA-98-3. Prepared by Tetra Tech, Inc., Owings Mills, MD and Maryland Department of Natural Resources, Monitoring and Non-Tidal Assessment Division. Annapolis, Maryland.

Stribling, J. B., S. R. Moulton, II, and G. Lester. 2003. Determining the quality of taxonomic data. *Journal of the North American Benthological Society* 22(4):621-631.

Suter, G. W. 1996. Abuse of hypothesis testing statistics in ecological risk assessment. *Human and Ecological Risk Assessment* 2:331-347.

Suter, II, G. W., S. B. Norton, and S. Cormier. 2002. A method for inferring the causes of observed impairments in aquatic ecosystems. *Environmental Toxicology and Chemistry* 21:1101-1111.

Taylor, B. R., and Z. Kovats. 1995. Review of artificial substrates for benthos sample collection. Prepared for Canada Centre for Mineral and Energy Technology (CANMET), Ottawa, Ontario, Canada.

Taylor, C. N., and C. E. Kuyatt. 1994. Guidelines for evaluating and expressing the uncertainty of NIST measurement results. NIST technical note 1297. National Institute of Standards and Technology, Department of Commerce, Washington, DC.

Taylor, J. K. 1988. Defining the accuracy, precision, and confidence limits of sample data. Pages 102-107 in L. H. Keith (editor). Principles of environmental sampling. ISBN 0-8412-1173-6. ACS professional reference book. American Chemical Society, Columbus, Ohio.

Thorp, J. H. 1992. Linkage between islands and benthos in the Ohio River, with implications for riverine management. Canadian Journal of Fisheries and Aquatic Sciences 49(9):1873-1882.

Thorp, J. H., and M. D. DeLong. 1994. The riverine productivity model: an heuristic view of carbon sources and organic processing in large river ecosystems. Oikos 70:305-308.

Tockner, K., and J. A. Stanford. 2002. Riverine flood plains: present state and future trends. Environmental Conservation 29:308-330.

Townsend, C. R., and A. G. Hildrew. 1976. Field experiments on the drifting, colonization and continuous redistribution of stream benthos. Journal of Animal Ecology 45:459-772.

Underwood, A. J. 1991. Beyond BACI: experimental designs for detecting human environmental impacts on temporal variations in natural populations. Australian Journal of Marine and Freshwater Research 42:569-587.

Underwood, A. J. 1992. Beyond BACI: the detection of environmental impacts on populations in the real, but variable world. Journal of Experimental Marine Biology and Ecology 161:145-178.

Underwood, A. J. 1994. On beyond BACI: sampling designs that might reliably detect environmental disturbances. Ecological Applications 4:3-15.

Urquhart, N. S., S. G. Paulsen, and D. P. Larsen. 1998. Monitoring for policy-relevant trends over time. Ecological Applications 8(2):246-257.

USEPA (US Environmental Protection Agency). 1990. Biological criteria: national program guidance for surface waters. EPA 440-5-90-004. Office of Water Regulations and Standards, US Environmental Protection Agency, Washington, DC.

USEPA (US Environmental Protection Agency). 1992. Procedures for initiating narrative biological criteria. EPA 822-B-92-002. Office of Science and Technology, US Environmental Protection Agency, Washington, DC.

USEPA (US Environmental Protection Agency). 1998. Lake and reservoir bioassessment and biocriteria technical guidance document. EPA 841-B-98-007. Office of Water, US Environmental Protection Agency, Washington, DC.

USEPA (US Environmental Protection Agency). 2000a. Stressor identification guidance document. EPA 822-B-00-025. US Environmental Protection Agency, Office of Water, Washington, DC.

USEPA (US Environmental Protection Agency). 2000b. Guidance for the data quality objectives process. EPA QA/G-4. EPA 600-R-96-055. US Environmental Protection Agency, Office of Environmental Information, Washington, DC.

USEPA (US Environmental Protection Agency). 2002. Summary of biological assessment programs and biocriteria development for states, tribes, territories, and interstate commissions: streams and wadeable rivers. EPA 822-R-02-048. US Environmental Protection Agency, Washington, DC. (available at: http://www.epa.gov/bioindicators/html/program_summary.html).

USEPA (US Environmental Protection Agency). 2005. Use of biological information to tier designated aquatic life uses in state and tribal water quality standards. EPA 822-R-05-001. Office of Water, US Environmental Protection Agency, Washington, DC.

USEPA (US Environmental Protection Agency). 2006. Technical guidance for assessing performance and comparability of bioassessment methods and their results. US Environmental Protection Agency, Office of Water, Washington, DC (in preparation).

Van Dam, H., A. Mertens, and J. Sinkeldam. 1994. A coded checklist and ecological indicator values of freshwater diatoms from the Netherlands. *Netherlands Journal of Aquatic Ecology* 28(1):117-133.

Vannote, R. L., G. W. Minshall, K. W. Cummins, J. R. Sedell, and C. E. Cushing. 1980. The river continuum concept. *Canadian Journal of Fisheries and Aquatic Sciences* 37:130-137.

Van Sickle, J. 1997. Using mean similarity dendograms to evaluate classifications. *Journal of Agricultural, Biological, and Environmental Statistics* 2:370-388.

Van Sickle, J., C. P. Hawkins, D. P. Larsen, and A. T. Herlihy. 2005. A null model for the expected macroinvertebrate assemblage in streams. *Journal of the North American Benthological Society* 24(1):178-191.

Van Sickle, J., D. D. Huff, and C. P. Hawkins. 2006. Selecting discriminant function models for predicting the expected taxa richness of aquatic macroinvertebrates. *Freshwater Biology* 51:359-372.

- Van Sickle, J., and R. M. Hughes. 2000. Classification strengths of ecoregion, catchments and geographic clusters for aquatic vertebrates in Oregon. *Journal of the North American Benthological Society* 19:370-384.
- Vincent, R. 1971. River electrofishing and fish population estimates. *Progressive Fish Culturist* 33(3):163–169.
- Vinson, M. R., and C. P. Hawkins. 1996. Effects of sampling area and subsampling procedure on comparisons of taxa richness among streams. *Journal of the North American Benthological Society* 15:392-399.
- Vitousek P. M., J. D. Aber, R. W. Howarth, G. E. Likens, P. A. Matson, D. W. Schindler, W. H. Schlesinger, and D. G. Tilman. 1997. Human alteration of the global nitrogen cycle: sources and consequences. *Ecological Applications* 7:737-750.
- Voelz, N. J., and J. V. McArthur. 2000. An exploration of factors influencing lotic insect species richness. *Biodiversity and Conservation* 9:1543-1570.
- Wade, D. C., and S. B. Stalcup. 1987. Assessment of the sport fishery potential for the bear creek floatway: biological integrity of representative sites, 1986. Report No. TVA/ONRED/AWR-87/30. Tennessee Valley Authority, Muscle Shoals, Alabama.
- Waite, I. R., A. T. Herlihy, D. P. Larsen, and D. J. Klemm. 2000. Comparing strengths of geographic and nongeographic classifications of stream benthic macroinvertebrates in the Mid-Atlantic highlands, USA. *Journal of the North American Benthological Society* 19:429-441.
- Wang, L., J. Lyons, and P. D. Kanehl. 1998. Development and evaluation of a habitat rating system for low-gradient Wisconsin streams. *North American Journal of Fisheries Management* 18(4):775-785.
- Wang, L., J. Lyons, P. D. Kanehl, and R. Gatti. 1997. Influences of watershed land use on habitat quality and biotic integrity in Wisconsin streams. *Fisheries* 22(6):6-12.
- Ward, J. V. 1989. Riverine-wetland interactions. Pages 385-400 in R. R. Sharitz and J. W. Gibbons (editors). *Freshwater Wetlands and Wildlife*. DOE Symposium Series No. 61, CONF-860301, USDOE Office of Scientific and Technical Information, Oak Ridge, Tennessee.
- Ward, J. V. 1998. Riverine landscapes: biodiversity patterns, disturbance regimes, and aquatic conservation. *Biological Conservation* 83:269-278.
- Ward, J. V., F. Malard, and K. Tockner. 2001. Landscape ecology: a framework for integrating pattern and process in river corridors. *Landscape Ecology* 17:35-45.
- Ward, J. V., and J. A. Stanford. 1983. The intermediate disturbance hypothesis: an explanation for biotic diversity patterns in lotic ecosystems. Pages 347-356 in T. D. Fontaine, III and S. M. Bartell (editors). *Dynamics of lotic ecosystems*. Ann Arbor Science, Ann Arbor, Michigan.

Ward, J. V., and J. A. Stanford. 1995. The serial discontinuity concept: extending the model to floodplain rivers. *Regulated Rivers Research and Management* 10:159-168.

Weber, C. I. (editor). 1973. Biological field and laboratory methods for measuring the quality of surface waters and effluents. EPA 670-4-73-001. Natural Environmental Research Center, Office of Research and Development, US Environmental Protection Agency Cincinnati, Ohio.

Wehr, J. D., and R. G. Sheath (editors). 2003. Freshwater algae of North America: ecology and classification. Academic Press, San Diego, California.

Welcomme, R. L. 1985. River fisheries. Food and Agriculture Organization Fisheries Technical Paper 262. Food and Agriculture Organization of the United Nations, Rome, Italy.

Welcomme, R. L. 1989. Floodplain fisheries management. Pages 209-233 in J. A. Gore, and G. E. Petts (editors). Alternatives in regulated river management. CRC Press, Boca Raton, Florida.

Welcomme, R. L., K. O. Winemiller, and I. G. Cowx. 2006. Fish environmental guilds as a tool for assessment of ecological condition of rivers. *River Research and Applications* 22:377-396.

Wetzel, R. G. 2001. Limnology: lake and river ecosystems. 3rd edition. Academic Press, New York.

Wetzel, R. L. 1979. Periphyton measurements and applications. In R. L. Wetzel (editor). Methods and measurements of periphyton communities. American Society of Testing and Material ASTM STP 690:3-33.

Whited, D. C., J. A. Stanford, and J. S. Kimball. 2003. Application of airborne multispectral digital imagery to characterize the riverine habitat. *Internationale Vereinigung für Theoretische und Angewandte Limnologie* 28(3):1373-1380.

Whittaker, R. H. 1952. A study of summer foliage insect communities in the Great Smoky Mountains. *Ecological Monographs* 22:1-44.

Wildhaber, M. L., P. J. Lamberson, and D. L. Galat. 2003. A comparison of measures of riverbed form for evaluating distributions of benthic fishes. *North American Journal of Fisheries Management* 23:543-557.

Wiens, J. A. 2002. Riverine landscapes: taking landscape ecology into the water. *Freshwater Biology* 47:501-515.

Wilhelm, J. G. O. 2002. A habitat rating system for non-wadeable rivers of Michigan. Master's Thesis, School of Natural Resources and Environment, University of Michigan, Ann Arbor, Michigan.

Wilhelm, J. G. O., J. D. Allan, K. J. Wessell, R. W. Merritt, and K. W. Cummins. 2005. Habitat assessment of non-wadeable rivers in Michigan. *Environmental Management* 36:592-609.

Wright, J. F. 1995. Development and use of a system for predicting the macroinvertebrate fauna in flowing waters. *Australian Journal of Ecology* 20:181-197.

Wright, J. F. 2000. An introduction to RIVPACS. Pages 1-24 in J. F. Wright, D. W. Sutcliffe, and M. T. Furse (editors). *Assessing the biological quality of freshwaters: RIVPACS and other techniques*. Freshwater Biological Association, Ambleside, Cumbria, UK.

Wright, J. F., M. T. Furse, and P. D. Armitage. 1993. RIVPACS: a technique for evaluating the biological water quality of rivers in the UK. *European Water Pollution Control* 3:15-25.

Wright, J. F., D. Moss, P. D. Armitage, and M. T. Furse. 1984. A preliminary classification of running-water sites in Great Britain based on macroinvertebrate species and the prediction of community type using environmental data. *Freshwater Biology* 14:221-256.

Wright, J. F., J. M. Winder, R. J. M. Gunn, J. H. Blackburn, K. L. Symes, and R. T. Clarke. 2000. Minor local effects of a River Thames power station on the macroinvertebrate fauna. *Regulated Rivers: Research and Management* 16(2):159-174.

Yin, Y., J. S. Winkelman, and H. A. Langrehr. 2000. Long term resource monitoring program procedures: aquatic vegetation monitoring. LTRMP Program Report 95-P002-7. US Geological Survey, Upper Midwest Environmental Sciences Center, La Crosse, Wisconsin.

Yoccoz, N. G. 1991. Use, overuse, and misuse of significance tests in evolutionary biology and ecology. *Bulletin of the Ecological Society of America* 72:106-111.

Yoder, C. O., R. J. Miltner, and D. White. 1999. Assessing the status of aquatic life designated uses in urban and suburban watersheds. In *Proceedings of the National Conference on Retrofit Opportunities for Water Resource Protection in Urban Environments*. EPA-625-R-99-002.

Yoder, C. O., and E. T. Rankin. 1995. Biological criteria program development and implementation in Ohio. Pages 109-144 in W. S. Davis and T. P Simon (editors). *Biological assessment and criteria: tools for water resource planning and decision making*. Lewis Publishers, Boca Raton, Florida.

Yoder, C. O., and M. A. Smith. 1999. Using fish assemblages in a state biological assessment and criteria program: essential concepts and considerations. Pages 17-56 in T. P. Simon (editor). *Assessing the sustainability and biological integrity of water resources using fish communities*. CRC Press, Boca Raton, Florida.

Zar, J. H. 1999. *Biostatistical Analysis*. 4th edition. Prentice Hall, Upper Saddle River, New Jersey.